

THE DARBOUX COORDINATES FOR A NEW FAMILY OF HAMILTONIAN OPERATORS AND LINEARIZATION OF ASSOCIATED EVOLUTION EQUATIONS

JIRINA VODOVÁ

ABSTRACT. A. de Sole, V. G. Kac, and M. Wakimoto have recently introduced a new family of compatible Hamiltonian operators of the form $H^{(N,0)} = D^2 \circ ((1/u) \circ D)^{2n} \circ D$, where $N = 2n + 3$, $n = 0, 1, 2, \dots$, u is the dependent variable and D is the total derivative with respect to the independent variable. We present a differential substitution that reduces any linear combination of these operators to an operator with constant coefficients and linearizes any evolution equation which is bi-Hamiltonian with respect to a pair of any nontrivial linear combinations of the operators $H^{(N,0)}$. We also give the Darboux coordinates for $H^{(N,0)}$ for any odd $N \geq 3$.

1. INTRODUCTION

The Hamiltonian evolution equations are well known to play an important role in modern mathematical physics. Indeed, a Hamiltonian operator maps the variational derivatives of the conserved quantities into symmetries; this plays an important role in the theory of integrable systems which often turn out to be *bi*-Hamiltonian, see e.g. [5, 6, 8, 13] and references therein. It is thus no wonder that the study and, in particular, the classification of Hamiltonian operators is a subject of ongoing interest, see for instance [1, 3, 7, 6, 11, 12] and the works cited there.

Recently A. de Sole, V. G. Kac, and M. Wakimoto have made a major advance in this area. Namely, in [4] they gave a conjectural classification of Poisson vertex algebras in one differential variable, i.e., of scalar Hamiltonian operators. *Inter alia*, they have come up with a new infinite family of compatible Hamiltonian operators $H^{(N,0)} = D^2 \circ ((1/u) \circ D)^{2n} \circ D$, where $N = 2n + 3$, $n = 0, 1, 2, \dots$, and D denotes the total derivative with respect to the space variable.

Although it is well known [15, 16] that there is no proper counterpart of the Darboux theorem on canonical forms of finite-dimensional Poisson structures for Hamiltonian operators associated with evolutionary PDEs, it is often possible to find new variables in which a Hamiltonian operator takes a simpler form. One such form is the Gardner operator D ; in analogy with the finite-dimensional case the associated new variables are often called the Darboux coordinates, see e.g. [3, 15]. Bringing a Hamiltonian operator into the Gardner form enables one e.g. to render the associated Hamiltonian systems into the canonical Hamiltonian form and construct Lagrangian representations (modulo potentialization) for these systems [16].

In this paper we present the transformations that bring the operators $H^{(N,0)}$ into the Gardner form, see Corollary 1 below. Moreover, in Theorem 1 we give a differential substitution which *simultaneously* turns the operators $H^{(N,0)}$ for all odd $N \geq 3$ into the operators with constant coefficients $\tilde{H}^{(N,0)} = -D^{2n+1}$.

2010 *Mathematics Subject Classification.* 37K05, 37K10.

Key words and phrases. Hamiltonian operators, bi-Hamiltonian systems, evolution equations, linearization.

These results could be employed e.g. for the study of (co)homology of the Poisson complexes associated with the operators $H^{(N,0)}$. The cohomologies in question play an important role e.g. in finding all Hamiltonian operators compatible with a given Hamiltonian operator and the associated multi-Hamiltonian systems, see for example [10, 14, 17] and references therein. Another possible application of the results in question is e.g. the construction of new integrable systems in spirit of [9] in the new variables from Theorem 1 or Corollary 1 with the subsequent pullback to the original variables. Last but not least, the differential substitution from Theorem 1 linearizes any evolution equation which is bi-Hamiltonian with respect to a pair of any nontrivial linear combinations of the operators $H^{(N,0)}$, thus exhibiting a broad class of somewhat unusual (in that they are C -integrable rather than S -integrable) integrable bi-Hamiltonian systems; see Corollary 2 for details.

2. PRELIMINARIES

In what follows we deal with Hamiltonian operators and associated Hamiltonian evolution equations involving a single spatial variable x and a single dependent variable u . An evolution equation of this kind has the form

$$u_t = K[u] = K(x, u, u_x, u_{xx}, \dots),$$

where the square brackets indicate that K is a *differential function* in the sense of [13], meaning that it depends on x , u , and finitely many derivatives of u with respect to the space variable x . Recall (see e.g. [5, 6, 13] for details) that an evolution equation is said to be *Hamiltonian* with respect to the Hamiltonian operator $\mathcal{D}$ if it can be written in the form

$$u_t = \mathcal{D}\delta_u \mathcal{T}[u],$$

where $\mathcal{T} = \int T[u]dx$ is the Hamiltonian functional, and δ_u denotes the variational derivative with respect to u .

Lemma 1 ([11]). *Let L_1 be a Hamiltonian operator in the variables x, u . Under the transformation*

$$x = \varphi(y, v, v_y, \dots, v_m), \quad u = \psi(y, v, v_y, \dots, v_n), \quad (1)$$

where $v_j = D_y^j(v)$, where D_y is the total derivative with respect to y , the operator L_1 goes into the Hamiltonian operator L_2 defined by the formula

$$\overline{L}_1 = (D_y(\varphi))^{-1} K^* \circ L_2 \circ K, \quad (2)$$

where $\overline{L}_1$ is obtained from L_1 under the substitution (1) and upon setting $D_x = (D_y(\varphi))^{-1} D_y$,

$$K = \sum_{i=0}^{\max(m,n)} (-1)^i D_y^i \circ \left(\frac{\partial \psi}{\partial v_i} D_y(\varphi) - \frac{\partial \varphi}{\partial v_i} D_y(\psi) \right),$$

and K^* is the formal adjoint of K .

Remark 1. Note that in general the operator L_2 may contain nonlocal terms unless (1) is a contact transformation, cf. e.g. [1, 4, 11].

3. THE MAIN RESULT

Theorem 1. *The transformation $x = v$, $u = 1/v_y$ turns the N th order Hamiltonian operator $H^{(N,0)} = D_x^2 \circ ((1/u) \circ D_x)^{2n} \circ D_x$, where $N = 2n + 3$, $n = 0, 1, 2, \dots$, into the Hamiltonian operator with constant coefficients $\tilde{H}^{(N,0)} = -D_y^{2n+1}$.*

The proof is obtained by a straightforward application of Lemma 1. Even though the transformation from Theorem 1 is not contact, in the particular case under study the transformed operators $\tilde{H}^{(N,0)}$ happen to be free of nonlocal terms (cf. Remark 1).

Using Lemma 1 we can further amplify the result of Theorem 1 by providing the Darboux coordinates (cf. Introduction) for the operators $H^{(N,0)}$.

Corollary 1. *The transformation $x = (-1)^{\frac{n+1}{2}} w_n$, $u = (-1)^{\frac{n+1}{2}} / w_{n+1}$, where $w_k = D_z^k(w)$, and z is the new independent variable, maps the Hamiltonian operator $H^{(N,0)} = D_x^2 \circ ((1/u) \circ D_x)^{2n} \circ D_x$ to the first-order Gardner operator D_z for any odd $N \geq 3$.*

Remark 2. The inverse of the transformation $x = v$, $u = 1/v_y$ is nothing but the extended hodograph transformation in the sense of [2]. Let us stress that, unlike the original transformation, the said inverse is not a differential substitution, i.e., it cannot be written in the form $y = \xi(x, u, u_x, \dots, u_q)$, $v = \chi(x, u, u_x, \dots, u_p)$.

In [4] the following conjecture is stated:

Conjecture 1 (De Sole, Kac, Wakimoto). *For any translation-invariant Hamiltonian operator H of order $N \geq 7$ there exists a contact transformation that brings H to either a quasiconstant coefficient skew-adjoint differential operator, or to a linear combination of the operators $H^{(j,0)}$ with $3 \leq j \leq N$, j odd.*

Recall that a differential function is called quasiconstant [4] if it depends only on x .

If we further allow for the transformation of the form $x = v$, $u = 1/v_y$ from Theorem 1 and use the latter, we can state a somewhat stronger conjecture:

Conjecture 2. *Any translation-invariant Hamiltonian operator H of order $N \geq 7$ can be transformed into a quasiconstant coefficient skew-adjoint differential operator using either a contact transformation or a composition thereof with the transformation $x = v$, $u = 1/v_y$.*

The transformation $x = v$, $u = 1/v_y$ can be written as a composition of the potentiation $x = z$, $u = w_z$ and of the hodograph transformation $z = v$, $w = y$. The latter is a contact (in fact, even a point) transformation, so, with the obvious change of notation, we can recast the above conjecture as follows:

Conjecture 3. *Any translation-invariant Hamiltonian operator H of order $N \geq 7$ can be transformed into a quasiconstant coefficient skew-adjoint differential operator using either a contact transformation or a composition thereof with the potentiation $x = y$, $u = v_y$.*

Thus, assuming that the above conjecture holds true, extending the class of allowed equivalence transformation using potentiation enables us to have just one type of normal forms for translation-invariant Hamiltonian operators, namely, the skew-adjoint differential operators with coefficients that depend only on x .

Upon further allowing for the transformations of the form $x = y, v = \sum_{i=0}^k b_i(x)v_i$ and using the following

Lemma 2. *Let $H = \sum_{i=0}^N a_i(x)D_x^i$ be quasiconstant skew-adjoint differential operator. There exists a transformation of the form $x = y, v = \sum_{i=0}^k b_i(x)v_i$ that turns H to the Gardner operator D_y .*

we arrive at a yet stronger conjecture:

Conjecture 4. *Any translation-invariant Hamiltonian operator of order $N \geq 7$ can be transformed into the Gardner operator D by either a transformation from Conjecture 3 or a composition thereof with a transformation of the form $x = y, v = \sum_{i=0}^k a_i(x)v_i$.*

4. APPLICATIONS TO EVOLUTION EQUATIONS

Recall that the transformation $x = v, u = 1/v_y$ can be written as the composition of the potentiation $x = z, u = w_z$ and of the hodograph transformation $z = v, w = y$. The first of these is nothing but introduction of the potential w for u . It is readily seen that a bi-Hamiltonian evolution equation

$$u_t = H_1 \delta_u \mathcal{T}_1 = H_2 \delta_u \mathcal{T}_2, \quad (3)$$

where $H_i = \sum_{j=1}^{k_i} c_{ij} H^{(N_{ij}, 0)}$, $i = 1, 2$, k_i are arbitrary natural numbers, and c_{ij} are arbitrary constants, is nothing but the pullback of the bi-Hamiltonian equation

$$w_t = \check{H}_1 \delta_w \check{\mathcal{T}}_1 = \check{H}_2 \delta_w \check{\mathcal{T}}_2, \quad (4)$$

where $\check{H}_i = \sum_{j=1}^{k_i} c_{ij} \check{H}^{(N_{ij}, 0)}$, $i = 1, 2$,

$$\check{H}^{(N, 0)} = -D_z \circ \left(\frac{1}{w_z} D_z \right)^{2n},$$

and $\check{\mathcal{T}}_i$ are obtained from $\mathcal{T}_i$ using the substitution $x = z, u = w_z$. It is natural to refer to (4) as to the potential form of (3).

Proposition 1. *The transformation $z = v, w = y$, where y is the new independent variable, linearizes the potential form (4) of the bi-Hamiltonian evolution equation (3).*

Before proving this let us point out the following important consequence of this result.

Corollary 2. *The differential substitution $x = v, u = 1/v_y$ relates any equation of the form (3) to a linear evolution equation with constant coefficients.*

Informally, this just means that the inverse of the transformation $x = v$, $u = 1/v_y$ linearizes (3), but this statement should be treated with some care, as this transformation is not uniquely invertible, and the inverse is not a differential substitution, cf. Remark 2 and [18].

Proof of Proposition 1. The hodograph transformation $z = v$, $w = y$ sends (4) into

$$v_t = \tilde{H}_1 \delta_v \tilde{\mathcal{T}}_1 = \tilde{H}_2 \delta_v \tilde{\mathcal{T}}_2,$$

where $\tilde{H}_i = \sum_{j=1}^{k_i} c_{ij} \tilde{H}^{(N_{ij},0)} = -\sum_{j=1}^{k_i} c_{ij} D_y^{N_{ij}-2}$ are linear differential operators with constant coefficients, and $\tilde{\mathcal{T}}_i$ are obtained from $\mathcal{T}_i$ using the transformation in question.

Lemma 3. *Let $v_t = K[v]$ be an n th order evolution equation which is bi-Hamiltonian with respect to a pair of Hamiltonian operators with constant coefficients. Then $v_t = K[v]$ is necessarily a linear equation with constant coefficients, i.e., we have $v_t = \sum_{i=0}^n c_i v_i$, where $c_i = \text{const}$.*

Proof of the lemma. Denote the Hamiltonian operators in question by $\mathcal{D}_i$, $i = 1, 2$. Since $v_t = K[v]$ is bi-Hamiltonian with respect to these operators by assumption, their ratio $\mathcal{R} = \mathcal{D}_2 \circ \mathcal{D}_1^{-1}$ is a (formal) recursion operator for this equation, that is (see e.g. [13] for details),

$$\text{pr } \mathbf{v}_K(\mathcal{R}) - [\mathcal{D}_K, \mathcal{R}] = 0,$$

where $\mathcal{D}_K = \sum_{i=0}^n (\partial K / \partial u_i) D^i$ is the Fréchet derivative of K and $\text{pr } \mathbf{v}_K$ is the prolongation of the evolutionary vector field $\mathbf{v}_K$ with the characteristic K . Since $\mathcal{D}_i$, $i = 1, 2$, have constant coefficients by assumption, we have $\text{pr } \mathbf{v}_K(\mathcal{D}_i) = 0$, and therefore $\text{pr } \mathbf{v}_K(\mathcal{R}) = 0$, so $\mathcal{D}_K$ commutes with $\mathcal{R}$, whence it readily follows that $\mathcal{D}_K$ has constant coefficients (recall that K is independent of t by assumption) and therefore K indeed is a linear combination of v_i with constant coefficients. $\square$

The desired result now readily follows from the above lemma. $\square$

Example 1. Consider a bi-Hamiltonian evolution equation

$$u_t = D_x^3 (u^{-2}) = H^{(3,0)} \delta_u \mathcal{T}_1 = H^{(5,0)} \delta_u \mathcal{T}_2, \quad (5)$$

where $\mathcal{T}_1 = -\int dx/u$ and $\mathcal{T}_2 = \int x^2 u dx$.

The potential form (4) of (5) reads

$$w_t = D_z^2 (w_z^{-2}) = \check{H}^{(3,0)} \delta_w \check{\mathcal{T}}_1 = \check{H}^{(5,0)} \delta_w \check{\mathcal{T}}_2. \quad (6)$$

Recall that $u = w_z$ and $x = z$; we have $\check{H}^{(3,0)} = -D_z$, $\check{H}^{(5,0)} = -D_z \circ ((1/w_z) D_z)^2$, $\check{\mathcal{T}}_1 = -\int dz/w_z$, and $\check{\mathcal{T}}_2 = \int z^2 w_z dz$. Note that (6) has, up to a rescaling of t , the form (2.31) from [2].

In perfect agreement with Proposition 1 (cf. also Proposition 2.2 in [2]) the hodograph transformation $z = v$, $w = y$ linearizes (6) into a (trivially) bi-Hamiltonian equation

$$v_t = -2v_{yyy} = \tilde{H}^{(3,0)} \delta_v \tilde{\mathcal{T}}_1 = \tilde{H}^{(5,0)} \delta_v \tilde{\mathcal{T}}_2, \quad (7)$$

where $\tilde{H}^{(3,0)} = -D_y$, $\tilde{H}^{(5,0)} = -D_y^3$, $\tilde{\mathcal{T}}_1 = -\int v_y^2 dy$, and $\tilde{\mathcal{T}}_2 = \int v^2 dy$.

The transformation $x = v$, $u = 1/v_y$ relates (7) to (5), cf. Corollary 2, so (5) provides an explicit example of a C -integrable (rather than S -integrable) bi-Hamiltonian system, just as discussed in Introduction.

ACKNOWLEDGEMENTS

The author thanks Dr. A. Sergyeyev for stimulating discussions. This research was supported in part by the Ministry of Education, Youth and Sports of the Czech Republic under grant MSM 4781305904.

REFERENCES

- [1] A. M. Astashov, A. M. Vinogradov, *On the structure of Hamiltonian operator in field theory*, J. Geom. and Phys. 3 (1986), no. 2, 263–287.
- [2] P. A. Clarkson, A. S. Fokas, M. J. Ablowitz, *Hodograph Transformations of Linearizable Partial Differential Equations*, SIAM J. Appl. Math. 49 (1989), 1188–1209.
- [3] D. B. Cooke, *Classification Results and the Darboux Theorem for Low-Order Hamiltonian Operators*, J. Math. Phys. 32 (1991), 109–119.
- [4] A. de Sole, V. G. Kac, M. Wakimoto, *On Classification of Poisson Vertex Algebras*, Transformation Groups (2010), to appear, arXiv:1004.5387.
- [5] L.A. Dickey, *Soliton Equations and Hamiltonian Systems*, World Scientific, River Edge, NJ, 2003.
- [6] I. Dorfman, *Dirac Structures and Integrability of Nonlinear Evolution Equations*, John Wiley and Sons, Chichester etc., 1993.
- [7] B. A. Dubrovin and S. P. Novikov, *Hydrodynamics of weakly deformed soliton lattices. Differential geometry and Hamiltonian theory* (Russian), Uspekhi Mat. Nauk 44, no.6, (1989), 29–98. English transl.: Russ. Math. Surv. 44 (1989) 35–124.
- [8] A. S. Fokas, I. M. Gel'fand, *Bi-Hamiltonian structures and integrability*, in: Important developments in soliton theory, Springer, Berlin, 1993, 259–282.
- [9] A. S. Fokas, P. J. Olver, P. Rosenau, *A plethora of integrable bi-Hamiltonian equations*, in: Algebraic aspects of integrable systems, Birkhäuser Boston, Boston, MA, 1997, 93–101.
- [10] I. Krasil'shchik, *Algebraic Theories of Brackets and Related (Co)Homologies*, Acta Appl. Math. 109 (2010), 137–150, arXiv:0812.4676.
- [11] O. I. Mokhov, *Hamiltonian differential operators and contact geometry* (Russian), Funkc. anal. i ego prilozh. 21 (1987), no.3, 53–60. English transl.: Funct. Anal. Appl. 21 (1987), 217–223.
- [12] O. I. Mokhov, *Symplectic and Poisson Geometry on Loop Spaces of Smooth Manifolds and Integrable Equations*, Harwood Academic Publishers, Amsterdam, 2001.
- [13] P. J. Olver, *Applications of Lie Groups to Differential Equations*, Springer, N.Y., 1993.
- [14] P. J. Olver, *BiHamiltonian Systems*, in: Ordinary and Partial Differential Equations, B.D. Sleeman and R.J. Jarvis, eds., Longman, N.Y. 1987, 176–193.
- [15] P. J. Olver, *Darboux' Theorem for Hamiltonian Differential Operators*, J. Diff. Equ. 71 (1988), 10–33.
- [16] P. J. Olver, *Dirac's Theory of Constraints in Field Theory and the Canonical Form of Hamiltonian Differential Operators*, J. Math. Phys. 27 (1986), 2495–2501.
- [17] A. Sergyeyev, *A Simple Way of Making a Hamiltonian System into a Bi-Hamiltonian One*, Acta Appl. Math. 83 (2004), 183–197, arXiv:nlin/0310012.
- [18] V. V. Sokolov, *Pseudosymmetries and Differential Substitutions* (Russian), Funkc. anal. i ego prilozh. 22 (1988), no.2, 47–56. English transl.: Funct. Anal. Appl. 22 (1988), 121–129.